



# Department of Defense Legacy Resource Management Program

PROJECT 03-198


## ***SPRNCA Water Needs Study Info for Decision-Makers***

Gretchen Kent / U.S. Army Garrison Fort Huachuca

This document is unclassified and may be released to the public.

Report Documentation Page			Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>2010</b>		2. REPORT TYPE		3. DATES COVERED <b>00-00-2010 to 00-00-2010</b>	
4. TITLE AND SUBTITLE <b>SPRNCA Water Needs Study Info for Decision-Makers</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>U.S. Army Garrison Fort Huachuca, Fort Huachuca, AZ, 85613</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>Same as Report (SAR)</b>	18. NUMBER OF PAGES <b>15</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			

# *SPRNCA Water Needs Study Info for Decision-Makers*



*San Pedro Riparian National Conservation Area – Water Needs Study*

# Background

**USPP Planning Goal:** Ensure an adequate GW supply to meet the reasonable needs of both the area's residents and property owners (current & future) and the SPRNCA

- Detailed reports at:  
<ftp://www-ftp.tucson.ars.ag.gov/rscott/>
- SOW jointly developed with USPP
- To be integrated, reviewed and published as a USGS report
- **Intent:** Provide information and tools to policy and decision-makers regarding the hydrologic requirements of the SPRNCA and potential management actions that may be taken to reduce the consumptive water uses within the SPRNCA without resulting in any negative effects on riparian resources






# Primary Objectives


- Determine the water needs of riparian vegetation, through the riparian growing season and throughout the SPRNCA to ensure its long-term ecological integrity
- Quantify the total water use of riparian vegetation within the SPRNCA
- Determine the source of water used by key riparian plant species within the SPRNCA



# Hydrological Monitoring & Analyses



## SPRNCA Boundary and the Riparian Corridor



Riparian Corridor was based on

1. Digital Elevation Model (DEM) contours at 6 - 9 meters above river bottom
2. Vegetation from 1997 NALC Land Cover Map and Joe Watts/Army Corps of Engrs Map 2000
3. Aerial photographs

# Classification of reaches into condition classes


## *Vegetation assessment model*

A model was developed that places sites into one of three condition classes, based on field collection of nine bioindicators (Veg. variables). Each bioindicator is sensitive to changes in SW or GW hydrology.

Each condition class is reflective of different levels of ecosystem functional capacity.

The assessment model can be used to track changes in the abundance of each condition class over time.

## *“State of the SPRNCA”*





# Hydrologic Characteristics of each Condition Class

<b>Cond. Class (CC)</b>	<b>Flow Permanence (1)</b>	<b>Dry Seasonal Max. Depth to GW (ft) (2)</b>	<b>GW Fluctuation (ft) (3)</b>	<b>Percent of SPRNCA (4)</b>
<b>1</b>	<b>Intermittent-Dry (<math>&lt; 60\%</math>)</b>	<b>Deep (<math>&gt; 11.5</math> ft)</b>	<b>Large (<math>&gt; 3.3</math> ft Diff. between monthly max and min)</b>	<b>9%</b>
<b>2</b>	<b>Intermittent-Wet (<math>60\%</math> to <math>95\%</math>)</b>	<b>Moderately Shallow (<math>8.2 - 11.5</math> ft)</b>	<b>Moderate (<math>1.7</math> to <math>3.3</math> ft)</b>	<b>49%</b>
<b>3</b>	<b>Perennial (<math>&gt; 95\%</math>)</b>	<b>Shallow (<math>&lt; 8.2</math> ft)</b>	<b>Small - Stable (<math>&lt; 1.7</math> ft difference)</b>	<b>38%</b>

(1) 0% = no flow the entire year, 100% = surface flow entire year

(2) Dry-season mean as averaged across the floodplain

(3) As averaged across the floodplain

(4) 4% of the SPRNCA has not yet been adequately sampled





## Condition Class 3 (wettest)

**38% of SPRNCA**



Perennial or near-perennial  
stream flow (present >95% of  
time)

Shallow ground-water (mean  
depth of <2.5 m across floodplain  
during dry season) with little  
seasonal fluctuation (<0.5 m/yr)

Tall, dense, multi-aged  
cottonwood-willow forests

Tamarisk subdominant or absent

Channel lined by dense  
herbaceous cover



## Cond. Class 2 ( intermediate )



**49% of SPRNCA**

**Stream flow present 60%-95% of time**

**Moderately deep and fluctuating ground water**

**Tamarisk has increased, although cottonwood-willow still dominant.**

**Streamside herbaceous cover is reduced, and mesic herb species have replaced hydric species.**



# Condition Class 1 (driest)

9% of SPRNCA



Stream flow present <60% of time

Deep (>3.5 m in dry season) and highly fluctuating (>1 m/yr) GW

Tamarisk dominant

Short shrublands with limited upper canopy cover

Sparse streamside herbaceous cover, dominated by mesic species such as bermuda grass



# 2003 Riparian Water Use – SV Subwatershed

## Water Use - Main stem from Intl. Border to Tombstone Gage for Riparian Corridor

Cover Type	Veg. Area (ac)	CU [acre-ft yr <sup>-1</sup> ]	2003 ET/unit area (ft)
Mesquite	1790-2400	4044-5443	2.3 *
Cottonwood/Willow (Perennial)	630	1981	3.2
Cottonwood/Willow (Intermittent)	290	389	1.3
Sacaton (< 3 m to groundwater)	280-410	486-718	1.7
Open Water	110	421	4.0
Salt Cedar	2.7 – 7.5	6-17	2.3
<b>Total</b>		<b>7328-8969</b>	
Corell et al. (1996)		7700 *	
Goodrich et al. (2000)		6590 **	

\*\* \* Ranges due to classification procedure employed by the Army COE.

All but Salt Cedar rounded to nearest 10 ac

\*\* \*\* Using baseflow information from Palominas, Charleston, and Tombstone Gages



# ***Variability in Riparian Water Use and Climate***

**High interannual variability in climate will translate to a similar amount of variability in the riparian GW use**




## ***Mesquite Water Use Rates per Unit Area***

<b>2001</b>	<b>2.2 ft/yr</b>
<b>2002</b>	<b>1.7 ft/yr</b>
<b>2003</b>	<b>2.3 ft/yr</b>


**Large flow events (like October 2000) also impact the system's hydrology. This event maintained a higher level of flow permanence for 12 to 18 months following the October 2000 events.**

# GIS-based Veg. Management and Riparian ET Tool



## **Example:**

Change all Mesquite in user-defined area to Sacaton  
(orange areas along the river)



## **Example:**

Evaluate the effect of a prescribed burn with user-supplied polygon map of burn areas

**Contact ARS to schedule a training session**



*San Pedro Riparian National Conservation Area – Water Needs Study*

# Study Period Conditions (Hydro-Climatic Context)


## **Charleston streamflow:**

- Lower than average in all seasons except Spring and Fall 2000.
- Streamflows generally decrease each study year.


## **Precipitation:**

- Summers above average at Coronado
- Summers below average at Tombstone

AVERAGE PRECIPITATION AT CORONADO NATL. MONUMENT



AVERAGE PRECIPITATION AT TOMBSTONE



# ***Major Hydro-climatic Conclusions***

---

- **The magnitude of regional ground-water contribution to the stream alluvium and evapotranspiration control seasonal ground-water variability and streamflow permanence**
- **October 2000-sized floods play a significant role in maintaining water in the system for 12 to 18 months following the event**